

EFFECT OF CARBAMAZEPINE (TEGRETOL) ON TESTICULAR PARAMETERS IN ADULT MALE MICE (*MUS MUSCULUS*)

Zainab Abd-Ali Sabr and Assit Prof. Zainab Karim

Department of Biology, College of Science for Women, University of Baghdad, Iraq

Emails: Zainb.b81@gmail.com, Zainabk_bio@csw.uobaghdad.edu.iq

Abstract:

The present study was aimed to determine the effects of different doses of Carbamazepine (CBZ) (Tegretol) on the organs testes and epididymis parameters of albino Swiss mice. The study sample included 24 male mice divided into 4 groups. Mice of groups 1, 2 and 3 were orally administrated with a daily dose 2.5, 5 and 10 mg/kg bw, Carbamazepine respectively, while group 3 served as control and administrated with 0.1 ml distilled water. The treatment continued for 60 days. Carbamazepine as a drug used in the treatment of epilepsy and neuropathic pain has been shown to stimulate the effects of free radicals and cause reproductive disorders. The present study investigates the effects of carbamazepine medication on serum sex hormones (Prolactin, LH, FSH and Testosterone) and some of sperm parameters such as (sperm count, motility, abnormality and viability). The results show that treatment with (CBZ) significant decrease ($p<0.05$) in sex hormones levels, also observed significant decreasing ($p<0.05$) in sperm count, sperm motility, when compared to that of normal control. The tests of the sperm morphology showed a number of deformations as a result of the treatment, such as the appearance of wavy tail, folded tail and amorphous head with hook tail. It can be concluded that sub-acute exposure to Carbamazepine exerts negative effects on testes and epididymis parameters of treated mice.

Key words: Carbamazepine , Sex hormones , Sperm parameters, Prolactin , Testes

Introduction:

Epilepsy is a chronic disease characterized by frequent bouts caused by a defect in the transmission of electrical signals in the brain, which affects about 50 million people worldwide(1,2). Carbamazepine ($C_{15}H_{12}N_2O$) is a tricyclic compound and commonly known as Tegretol and one of the most famed drugs used to treat epilepsy. Carbamazepine is an anticonvulsant and mood-stabilizing drug used mainly in the management of bipolar disorder (trigeminal neuralgia), attention-deficit hyperactivity disorder (ADHD), schizophrenia, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, disorder, borderline, and post-traumatic stress disorder(3,4). Carbamazepine is a sodium channel blocker. It binds preferentially to voltage-gated sodium channels in their inactive conformation, which prevents repetitive and sustained firing of an action potential. Carbamazepine has effects on serotonin systems but the relevance to its antiseizure effects is uncertain(5). Large studies showed the reproductive hormonal and sexual dysfunction occur in men with epilepsy(28-70 %) and showed as decreased libido, infertility and erectile difficulties(6,7). CBZ induce the metabolism of sex steroid hormones and causes of reproductive and sexual dysfunction.(6, 8).

Materials and Methods:

Preparation of carbamazepine required dosage:

Dose of carbamazepine was prepared depending on the half lethal dose (LD_{50}) of the drug which was 529 mg/kg b.w.(9). One dose of carbamazepine was selected 200mg/kg to test its toxic effect, while the weights of mice used in the experiments was ranged between 25-30g. The required dose of the drug was dosed orally for all mice once per day for 60 days for three concentrations (2.5 , 5, 10 mg/kg) . It was possible to calculate the amount of drug used for the mice in this study based on the following equation.

$$X/D = W/1000$$

Experiment animals:

Adult white male mice used in this study (24 animals) with an average weight of about (25-30 g), were used in this study, obtained from the Animal House of the Iraqi center for Cancer Research. The animals were placed in plastic cages and their floors were covered with sawdust. The cages were cleaned in terms of replacing sawdust twice during the week, and the cages were sterilized with disinfectants to avoid diseases. During the experiment stages, all animals were subjected to similar laboratory conditions in terms of continuous ventilation and lighting (12 hours of light and 12 hours of darkness), and the temperature ranged between (22-25 ° C) and humidity between (40-60%). Water and suspension were given. Towards continuous duration of the study.

Experiment groups:

The animals were divided randomly into four groups (each group contains 6 animals). the animals were given Carbamazepine with concentrations of (2.5, 5 ,10 mg/kg). Administrated as aqueous solution of the substance orally .The fourth group is control, were give D.W .

Blood collection:

Blood was collected in one stage from the animals by a heart stab method while they were alive. The blood was kept in anticoagulant tubes (EDTA tube), then it was placed in a cooled centrifuge at a mouse of 3000 rpm / minute at a temperature of 25°C for a period of 10 minutes, in order to obtain the serum, after isolating the serum in sterile tubes, tests were performed to see Levels of the following hormones (Testosterone , Prolactin , LH, FSH) were done at the Biotechnology Research Center / Al-Nahrain University.

The total sperms count

For purpose of studying the specifications of the sperms , The tail of epididymis was taken and embedded in one drop of phosphate buffer solution at 37 °C in a watch glass , then the tail was cut in to at least 200 section by microsurgical scissors, in order to calculate sperm-specific parameters including , Motility, Viability , and then added one drop of Eosin-Nigrosine stain and mixed. Random fields of the sperm preparation were microscopically examined to assessed the percentage of sperms with abnormalities such as changes in head and tail , through the equation below:

$$\text{Percentage of abnormal sperms} = \frac{\text{Number of abnormal sperms}}{\text{total number}} \times 100$$

Statistical Analysis:

The Statistical Analysis System- SAS (2012) program was used to detect the effect of difference factors in study parameters. Least significant difference -LSD test (Analysis of Variation-ANOVA) was used to significant compare between means in this study (10).

Results:

The main results are shown in tables:

Table (1): The effect of different concentrations of the carbamazepine on the levels of prolactin , FSH , LH and testosterone

Groups	Mean ± SE			
	FSH (m IU\ml)	LH (m IU\ml)	Testosterone (Ng\ml)	Prolactin (m IU\ml)
Control	432.38 ± 75.88 a	139.93 ± 18.87 a	3.49 ± 0.45 a	14.39 ± 1.19 a
CBZ 2.5 mg\kg	265.34 ± 30.47 b	47.34 ± 4.23 b	1.85 ± 0.32 bc	9.09 ± 0.92 b
CBZ 5 mg\kg	244.59 26.76 b	47.85 ± 5.17 b	2.12 ± 0.37 b	8.95 ± 0.63 b
CBZ 10mg\kg	131.55 ± 12.38 b	34.80 ± 5.10 b	0.83 ± 0.12 c	4.55 ± 0.49 c
LSD value	133.93 *	31.83 *	1.053 *	2.64 *
Means having with the different letters in same column differed significantly. * (P≤0.05) .				

The results also showed a significant decrease (P≤0.05) in concentration FSH in groups treated with a drug for concentrations (2.5 ,5 ,10) mg\kg of body weight (265.34 ± 30.47) (244.59 ± 26.76) (131.55 ± 12.38) international unite respectively compared with the control group (432,38 ± 75.88) IU.As for the hormone LH, the results showed a significant decrease (P≤0.05) in the levels of the hormone in the tree groups treated with drug for concentrations (2.5 ,5 ,10) mg\kg of body weight (47.34 ± 4.23) (47.85 ±5.17) (34.80 ± 5.10) IU compared with the control group (139.93 ± 18.87) IU.The results obtained found significant decrease (P≤0.05) in the levels of testosterone hormone in the three groups treated with CBZ for concentrations (2.5,5 ,10) mg\kg of body weight . (1.85 ± 0.32) (2.12 ± 0.37) (0.83 ± 0.12) ng\ml compared with the control group (3.49 ± 0.45) ng\ml.The results also showed a significant decrease (P≤0.05) in the levels prolactin hormone in groups treated with CBZ for concentrations (2.5, 5, 10) mg\kg of body weight (9.09 ± 0.92) (8.95 ± 0.63) (4.55 ± 0.49) IU respectively, compared with the control group (14.39 ± 1.19) IU as in table (1).

The results showed found high significant decrease (P≤0.05) in total sperm count in group treated with drug for concentrations (2.5,5,10) mg/kg of body weight (20.25 ±2.01) (13.50 ±1.04) (11.00 ±0.91) sperm X106 compared with control group (28.50 ±1.84) sperm X106. As for the

motility of the sperm ,the results showed found significant decrease ($P \leq 0.05$) in sperm motility in group treated for concentrations (2.5, 5, 10) mg/kg of body weight (85.00 ± 2.04) (73.75 ± 4.26) (58.75 ± 4.26)% compared with control group (91.75 ± 1.37)%.

The results obtained the presence of abnormal forms of sperm in group treated for concentrations (2.5,5,10) mg/kg of body weight. (13.50 ± 1.55)(14.50 ± 1.32) (22.25 ± 2.01)% compared with control group (8.75 ± 1.49)%. The results showed a significant increase ($P \leq 0.05$) in the percentage of dead sperm in the groups treated for concentrations (2.5,5,10) mg/kg of body weight (12.2 ± 1.25) (19.25 ± 1.11) (25.50 ± 1.84) % respectively compared with the control group (8.0 ± 1.29)% as in table (2) .

Table (2): The effect of different concentrations of the carbamazepine on total count sperm ,motility, abnormality and viability 10^6

Group	Mean \pm SE			
	Total count sperm $\times 10^6$	Motility %	Abnormality %	Viability %
Control	28.50 \pm 1.84 a	91.75 \pm 1.37 a	8.75 \pm 1.49 c	8.00 \pm 1.29 c
CBZ 2.5mg/kg	20.25 \pm 2.01 b	85.00 \pm 2.04 a	13.50 \pm 1.55 bc	12.25 \pm 1.25 c
CBZ 5 mg/kg	13.50 \pm 1.04 c	73.75 \pm 4.26 b	14.50 \pm 1.32 b	19.25 \pm 1.11 b
CBZ 10 mg/kg	11.00 \pm 0.91 c	58.75 \pm 4.26 c	22.25 \pm 2.01 a	25.50 \pm 1.84 a
LSD value	4.722 *	10.046 *	4.982 *	4.323 *

Means having with the different letters in same column differed significantly. * ($P \leq 0.05$).

The results showed appearance of deformed sperms , including cytoplasmic droplet (Figure 2) , amorphous head with hook tail (Figure 3) , wavy tail (figure 4), folded tail (figure 5) , atrophy head and folded tail (figure 6) , headless sperm (figure 7) , bended tail (figure 8) , compare with the normal shape of sperm (figure 1) .




Fig 1 . Normal sperm of mice



Fig 2 . Cytoplasmic droplet



Fig. 3: Amorphous head with hook tail




Fig. 4: Wavy tail



Fig. 5: Folded tail




Fig.6: Atrophy head with folded tail



Fig. 7: Headless sperm



Fig. 8: Bended tail

Discussion:

The results of the present study indicated that a change occurred in serum sex hormones balance and significant decrease were showed in sperm count, sperm motility during CBZ. CBZ and some chemicals induce the metabolism of sex steroid hormones resulting in reduced blood concentration of bioactive testosterone, which results in diminished potency(8,11). The plasma concentration of free testosterone is decreased in male epileptic patients and same trend was observed in rats treated with anti-epileptic drugs and used androgen (12,13,14). This suggests that CBZ curb fertility in rats, not only by interfering with testicular functions, but maybe by affecting the hypothalamo-pituitary-testicular (gonadal) axis and thus decreasing the amount of testosterone in circulation. Testicular functions and male reproductive system are almost completely controlled by testosterone (15). The anti-epileptic drugs act on the pituitary gland and decline the secretion of FSH and LH responsible for spermatogenesis. The process of spermatogenesis and accessory reproductive organs are androgen dependent. Produced androgen production will cause a decrease in the number of mature Leydig cells and their functional cases.(16,17).

The present study found a significantly lower in prolactin hormone levels during CBZ-treated male mice than control group. The evidence shows that single AEDs can alter the responsiveness of the anterior pituitary to various stimuli (18).

Norepinephrine (NE) may stimulate LH release and serotonin(5-HT) may have an inhibitory and/or a stimulatory effect. The release of PRL from the anterior pituitary is primarily controlled by an inhibitory influence mediated by the tuberoinfundibular dopaminergic pathway (19).

The results of the present study indicated that a decrease in sperm count and showed decrease in sperm motility during CBZ medication. Also our study showed the presence of abnormal forms of sperm and obtained increase in the percentage of dead sperm. AED can directly act on the HPG axis or on peripheral hormones and induce various reproductive or sex hormonal abnormalities (20) AED has been shown to decrease sperm quantity and quality ,also affect spermatogenesis, resulting in infertility (21,22). Disturbances in reproductive hormones and gonadal morphology have been observed in patients with epilepsy and in non-epileptic animals on AED (23,24). Studies on AED including CBZ (25,26) indicated drug-related side effects which include abnormal sperm morphology , decreased motility, lower sperm count , it has been suggested that semen quality may be harmfully affected as (1) a result of direct toxic effect of AED on the gonads(27). (2)a side effect of AEDs on the HPG axis with abnormalities of sex hormones (such as testosterone) which are essential for normal sperm (28). (3) a result of carnitine deficiency associated with some AED including CBZ (29,30,31).

L-carnitine is produced by hepatocytes, concentrated in the blood stream, secreted by epididymal epithelium and taken up by the spermatozoa . L-carnitine and acetyl-carnitine play a crucial role in sperm metabolism, maturation and motility . Low levels of carnitine decline fatty acid concentrations within the mitochondria leading to reduce energy production and decrease sperm motility (32). CBZ was found to result in germ cell mutagens and an increase in the number of sperm with head defects in experimental studies (33). The severity in the decreasing of sperm counts and motility is drug dependent. That is because to direct effect CBZ on spermatogenesis,

specifically its interference with acetylcholine-esterase and glucose, both of which are required for the process of spermatogenesis. That is because sperm motility, in addition to being under the control of acetylcholine (34). also relies on glucose availability. Enzyme induction can also contribute to the development of chronic harmful effects, such as decreased sexual dysfunction, and potentially deleterious changes in cholesterol concentrations and other markers of vascular risk (35). CBZ-evoked oxidative harm to the spermatozoa could be as a result of excessive proteins in the epididymis and testis , androgen decreased causing sperm abnormality (36,37) . The elevated reactive oxygen species (ROS) that may harm sperm membrane causing changing morphology (38), also directs effect of AEDs and some drugs on spermatozoa germinal lineage cells because these drugs are highly lipid-soluble and cross the blood-testis barrier in to the seminiferous tubules. (39,40)

The reduction in the percentage viability of sperm could be interconnected with the reduction in sperm motility, since non-motile or slow spermatozoa were usually considered dead and could not be counted as live spermatozoa (41).

Conclusion:

The present study has shown that administration of CBZ (2.5,5,10) mg/kg to male mice, reduced serum sex hormones levels. our study indicate that decrease in sperm count and motility, presence of abnormal forms of sperm and increase of percentage of dead sperm .This study showed statistical significant changes when compared to control group.

References:

- 1: Engel Jr J. A proposed diagnostic scheme for people with epileptic seizures and with epilepsy: report of the ILAE Task Force on Classification and Terminology. *Epilepsia*. 2001 Jun;42(6):796-803.
2. Grzesiak AL, Lang M, Kim K, Matzger AJ. Comparison of the four anhydrous polymorphs of carbamazepine and the crystal structure of form I. *Journal of pharmaceutical sciences*. 2003 Nov 1;92(11):2260-71.
3. Kramer MA, Cash SS. Epilepsy as a disorder of cortical network organization. *The Neuroscientist*. 2012 Aug;18(4):360-72.
4. Rho JM, Sankar R. The pharmacologic basis of antiepileptic drug action. *Epilepsia*. 1999 Nov;40(11):1471-83.
5. Granger P, Biton B, Faure C, Vige X, Depoortere H, Graham D, Langer SZ, Scatton B, Avenet P. Modulation of the gamma-aminobutyric acid type A receptor by the antiepileptic drugs carbamazepine and phenytoin. *Molecular pharmacology*. 1995 Jun 1;47(6):1189-96.
- 6: Bauer J, Blumenthal S, Reuber M, Stoffel-Wagner B. Epilepsy syndrome, focus location, and treatment choice affect testicular function in men with epilepsy. *Neurology*. 2004 Jan 27;62(2):243-6.
- 7: Isojarvi JI, Repo M, Pakarinen AJ, Lukkarinen O, Myllylä VV. Carbamazepine, phenytoin, sex hormones, and sexual function in men with epilepsy. *Epilepsia*. 1995 Apr;36(4):366-70.

- 8: Hamed S, Mohamed K, El-Taher A, Hamed E, Omar H. The sexual and reproductive health in men with generalized epilepsy: a multidisciplinary evaluation. International journal of impotence research. 2006 May;18(3):287-95.
9. Abd Ali IK, Ibrahim TA, Farhan AD, Abbas MN. Study of the effect of pesticide 2, 4-D on the histological structure of the lungs in the albino mice (*Mus musculus*). Journal of Pharmaceutical Sciences and Research. 2018 Jun 1;10(6):1418-21.
10. Cary N. Statistical analysis system, User's guide. Statistical. Version 9. SAS. Inst. Inc. USA. 2012.
11. Rahman SA, Sattar DA. Effect of different concentration of Super Cyren pesticide on some physiological and histological traits of mice after different periods of oral administration. Iraqi Journal of Science. 2017;2291-300.
12. Verrotti A, Loiacono G, Laus M, Coppola G, Chiarelli F, Tiboni GM. Hormonal and reproductive disturbances in epileptic male patients: emerging issues. Reproductive Toxicology. 2011 May 1;31(4):519-27.
13. Osuntokun OS, Olayiwola G, Oladele A, Ola I, AyokaAbiodun O. Chronic administration of gabapentin and a gabapentin-carbamazepine combination reversibly suppress testicular function in male Wistar rats (*Rattus norvegicus*). Pathophysiology. 2017 Jun 1;24(2):63-9
14. Hassan EA, Hameed RY, Mezil SA. Impacts of Androgen Abuse and Overtraining on Endocrine Profile in Bodybuilders. Iraqi Journal of Science. 2021 Dec 30:4614-20.
15. Daoud AS, Bataineh H, Otoom S, Abdul-Zahra E. The effect of Vigabatrin, Lamotrigine and Gabapentin on the fertility, weights, sex hormones and biochemical profiles of male rats. Neuroendocrinology Letters. 2004 Jun 1;25(3):178-83.
16. Otoom S, Batieneh H, Hassan Z, Daoud A. Effects of long-term use Topiramate on fertility and growth parameter in adult male rats. Neuroendocrinology Letters. 2004 Oct 1;25(5):351-5.
17. Yenzeel JH. Effect of body mass index on sperm parameters and sex hormone level in sample of infertile iraqi men. Iraqi Journal of Science. 2017:1428-36.
18. Murialdo G, Manni R, De Maria A, Bonura ML, Polleri A, Tartara A. Luteinizing hormone pulsatile secretion and pituitary response to gonadotropin releasing hormone and to thyrotropin releasing hormone in male epileptic subjects on chronic phenobarbital treatment. Journal of endocrinological investigation. 1987 Feb;10(1):27-31.
19. Clemens JA. Monoaminergic control of the neuroendocrine system. InNeuroregulation of Autonomic, Endocrine and Immune Systems 1986 (pp. 243-264). Springer, Boston, MA.
20. Toone BK, Wheeler M, Nanjee M, Fenwick P, Grant R. Sex hormones, sexual activity and plasma anticonvulsant levels in male epileptics. Journal of Neurology, Neurosurgery & Psychiatry. 1983 Sep 1;46(9):824-6.
21. Taneja N, Kucherla K, Jain S, Maheshwari MC. Effect of phenytoin on semen. Epilepsia. 1994 Jan;35(1):136-40.
22. Xiaotian X, Hengzhong Z, Yao X, Zhipan Z, Daoliang X, Yumei W. Effects of antiepileptic drugs on reproductive endocrine function, sexual function and sperm parameters in Chinese Han men with epilepsy. Journal of Clinical Neuroscience. 2013 Nov 1;20(11):1492-7.

23. RØste LS, Taubøll E, Berner A, Berg KA, Aleksandersen M, Gjerstad L. Morphological changes in the testis after long-term valproate treatment in male Wistar rats. *Seizure.* 2001 Dec 1;10(8):559-65.
24. RØste LS, Taubøll E, Isojärvi JI, Berner A, Berg KA, Pakarinen AJ, T HUHTANIEMI IL, Knip M, Gjerstad L. Gonadal morphology and sex hormones in male and female Wistar rats after long-term lamotrigine treatment. *Seizure.* 2003 Dec 1;12(8):621-7.
25. RØste LS, Taubøll E, Haugen TB, Bjørnenak T, Saetre ER, Gjerstad L. Alterations in semen parameters in men with epilepsy treated with valproate or carbamazepine monotherapy. *European Journal of Neurology.* 2003 Sep;10(5):501-6.
26. Hayashi T, Yoshinaga A, Ohno R, Ishii N, Kamata S, Watanabe T, Yamada T. Asthenozoospermia: Possible association with long-term exposure to an anti-epileptic drug of carbamazepine. *International journal of urology.* 2005 Jan;12(1):113-4.
27. RØste LS, Taubøll E, Berner A, Berg KA, Aleksandersen M, Gjerstad L. Morphological changes in the testis after long-term valproate treatment in male Wistar rats. *Seizure.* 2001 Dec 1;10(8):559-65.
28. Rajender S, Monica MG, Walter L, Agarwal A. Thyroid, spermatogenesis, and male infertility. *Front Biosci (Elite Ed).* 2011 Jun 1;3:843-55.
29. Hamed SA, Abdella MM. The risk of asymptomatic hyperammonemia in children with idiopathic epilepsy treated with valproate: relationship to blood carnitine status. *Epilepsy research.* 2009 Sep 1;86(1):32-41.
30. Coppola G, Epifanio G, Auricchio G, Federico RR, Resicato G, Pascotto A. Plasma free carnitine in epilepsy children, adolescents and young adults treated with old and new antiepileptic drugs with or without ketogenic diet. *Brain and Development.* 2006 Jul 1;28(6):358-65.
31. Hug G, McGraw CA, Bates SR, Landrigan EA. Reduction of serum carnitine concentrations during anticonvulsant therapy with phenobarbital, valproic acid, phenytoin, and carbamazepine in children. *The Journal of pediatrics.* 1991 Nov 1;119(5):799-802.
32. Jeulin C, Lewin LM. Role of free L-carnitine and acetyl-L-carnitine in post-gonadal maturation of mammalian spermatozoa. *Human Reproduction Update.* 1996 Mar 1;2(2):87-102.
33. Shetty AJ. The effect of gabapentin and phenytoin on sperm morphology in Wistar rats. *Reprod Biol.* 2007 Nov 1;7(3):247-51.
34. Nelson L. Quantitative evaluation of sperm motility control mechanisms. *Biology of Reproduction.* 1972 Apr 1;6(2):319-24.
35. Perucca E.;1. Clinically relevant drug interactions with antiepileptic drugs. *British Journal of Clinical Pharmacology.* 2005(61):246–255.
36. Rao MV, Chinoy NJ. Effect of estradiol benzoate on reproductive organs and fertility in the male rat. *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 1983 Jul 1;15(3):189-98.
37. Akorede GJ, Ambali SF, Olatunji AO, Shittu M, Aremu A, Basiru A, Biobaku KT, Ahmed AO, Ameen SA. Protective effect of vitamin C on chronic carbamazepine-induced reproductive toxicity in male wistar rats. *Toxicology reports.* 2020 Jan 1;7:269-76.

38. Osuntokun OS, Olayiwola G, Oladele A, Ola I, AyokaAbiodun O. Chronic administration of gabapentin and a gabapentin-carbamazepine combination reversibly suppress testicular function in male Wistar rats (*Rattus norvegicus*). *Pathophysiology*. 2017 Jun 1;24(2):63-9.
39. Al-Timimi ZK. METRONIDAZOLE INDUCES SIGNIFICANT PATHOLOGICAL ALTERATIONS IN THE MALE REPRODUCTIVE SYSTEM OF MICE. *IRAQI JOURNAL OF AGRICULTURAL SCIENCES*. 2021 Dec 22;52(6):1375-81.
40. OKUMURA K. PERMEABILITY OF DRUGS AND CHEMICALS ACROSS THE BLOOD-TESTIS BARRIER. *Journal of toxicological sciences*. 1982 Nov 25;7(4):312..
41. Raji Y., Udoh U.S., Mewoyeka O.O., Ononye F.C. and Bolarinwa A.F..Implication of reproductive endocrine malfunction in male antifertility efficacy of *Azadirachtaindica* extractin rats. *African journal of Medicine and Medical Sciences*. 2007(32): 159-165.